

N69-10810

RESULTS OF AIRBORNE SURVEY OF NO₂ AND SO₂
OVER LOS ANGELES BASIN

Barringer Research Limited
Rexdale, Ontario, Canada

September 1968

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

NG9-10810

RESULTS OF AIRBORNE SURVEY
OF NO₂ AND SO₂ OVER
LOS ANGELES BASIN

PREPARED FOR
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
MANNED SPACECRAFT CENTER, HOUSTON
AND
THE UNITED STATES GEOLOGICAL SURVEY
IN ASSOCIATION WITH
THE GEOGRAPHY DEPARTMENT OF THE UNIVERSITY OF CALIFORNIA
CONTRACT NO. NAS9-7958

PREPARED BY
BARRINGER RESEARCH LIMITED
304 Carlingview Drive
Rexdale, Ontario, Canada

TR-68-68

SEPTEMBER - 1968

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

NOTICE

THIS DOCUMENT HAS BEEN REPRODUCED FROM THE BEST COPY FURNISHED US BY THE SPONSORING AGENCY. ALTHOUGH IT IS RECOGNIZED THAT CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED IN THE INTEREST OF MAKING AVAILABLE AS MUCH INFORMATION AS POSSIBLE.

TABLE OF CONTENTS

<u>SECTION</u>	<u>DESCRIPTION</u>	<u>PAGE NO.</u>
1	General & Background	1
2	Flight Patterns & Schedules	2
3	Data Reduction	3
4	Discussion of Results	5
5	Comparison of Aircraft with Surface Data,	8
6	Conclusions	9

LIST OF TABLES

TABLE

1	Summary of Flights	
2	Data Reduction	Flight 2
3	Temperature Sounding	Flight 2
4	Temp/Press Profile Chart	Flight 2
5	Data Reduction	Flight 3
6	Temperature Sounding	Flight 3
7	Temp/Press. Profile Chart	Flight 3
8	Temp/Press Profile Chart	Flight 3
9	Temp/Press Profile Chart	Flight 3
10	Data Reduction	Flight 4
11	Temperature Sounding	Flight 4
12	Temp/Press Profile Chart	Flight 4
13	Data Reduction	Flight 5
14	Temperature Sounding	Flight 5
15	Temp/Press Profile Chart	Flight 5
16	Data Reduction	Flight 6
17	Temperature Sounding	Flight 6
18	Temp/Press Profile Chart	Flight 6
19	Comparison Between Aircraft and Ground NO ₂ Data	

LIST OF FIGURES

FIGURE

COMPARISON BETWEEN AVERAGE
POLLUTION and SURFACE POLLUTION

LIST OF APPENDICES

APPENDIX

A	Derivation of Pathlength Factor
B	Calculation of Concentration from Total Vertical Gas Burden
C	Ground Truth Air Pollution Data Acquired by Los Angeles Air Pollution Control District
D	Weather Data for the Period May 21st to May 24th, 1968.

LIST OF DRAWINGS

CHART

I	Airborne NO ₂ Survey	Flight 2
II	Airborne SO ₂ Survey	Flight 3
III	Airborne NO ₂ Survey	Flight 4
IV	Airborne SO ₂ Survey	Flight 5
V	Airborne NO ₂ Survey	Flight 6

1. GENERAL AND BACKGROUND

An airborne survey of atmospheric nitrogen dioxide (NO_2) and sulfur dioxide (SO_2) was performed over the Los Angeles Basin including Los Angeles and Riverside Counties. The purpose of the survey was to evaluate the Barringer correlation technique as a means of remotely measuring concentrations of atmospheric contaminants. This work formed one segment of a large scale multi-sensor testing program managed and directed by the United States Geological Survey in association with the Geography Department of the University of California, Riverside.

The objectives of the larger program was to test the utility of various remote sensing techniques for space applications as related to the Geography discipline of the U.S.G.S./N.A.S.A. Earth Resources Program. Two contrasting areas in southern California were designated for intensive multi-sensor testing during the period May 21st to June 4th inclusive namely (1) the city of Los Angeles showing a cross section of residential developments of wide range and quality, industrial areas, portions of transportation network, commercial districts and air pollution and (2) a concentration of irrigation, agriculture and small towns in the Coachella and Imperial Valleys in the Salton Sea portion of the Colorado Desert, about 100 miles from coastal Los Angeles.

The work reported on herein is confined to air pollution measurements in the Los Angeles Basin over the period May 21st to May 24th inclusive. The aircraft used for pollution sensing was an Aero Commander 500A based at Riverside Municipal Airport.

This report describes the airborne measurement procedures, the flight lines or survey patterns flown, the meteorological conditions at the time of flight, the method of data reduction and the end results obtained. These results are compared with simultaneous ground based measurements provided by A.P.C.D. (Air Pollution Control District) stations.

2. FLIGHT PATTERNS AND SCHEDULES

Flight patterns were chosen to provide a general picture of atmospheric pollution over the greater Los Angeles area encompassing the western portion of Riverside County, the San Fernando Valley and the metropolitan Los Angeles area south to Long Beach. Consideration was also given to the selection of each leg or line of the patterns to obtain pollutant profiles which would overlay A.P.C.D. stations and hence provide most meaningful comparisons of ground level pollutant concentrations. For the last day of measurements, Friday May 24th, data runs were made for SO_2 and NO_2 along the same north-south and east-west flight lines used by the N.A.S.A. 240A (flown on the same day) aircraft. Tables 1 and 2 summarize the flight numbers, flight durations and cloud conditions. Useful survey time was limited by heavy cloud formation over the target area particularly on the first day, Tuesday, May 21st when rapid cloud build up forced termination of the flight after an initial calibration and before much useful data could be obtained. Meteorological conditions improved gradually from Tuesday May 21st to Friday May 24th with the best conditions on Thursday and Friday afternoons (flights 3, 4, 5 and 6). The flight patterns and pollutant profiles are shown in Figures 2, 3, 4 and 5 of the report.

As a rule the procedure followed in accomplishing each flight or mission was to obtain a vertical temperature profile of the atmosphere by monitoring ambient air temperature at regular intervals during a spiral ascent over Riverside Airport. The operational altitude for the survey was chosen on the basis of the temperature inversions indicated by these data and by visual observations of haze layer altitudes. In general the aircraft altitude was significantly higher than the inversion altitude. A spectrometer calibration was then performed before commencing a data run over the first leg of the pattern.

This consisted of zeroing the spectrometer output against the overhead sky and then recording the response obtained by temporarily inserting a standard reference cell in the optical path of the instrument. A similar calibration was performed frequently during the mission generally at the beginning and/or end of each leg of the flight patterns. At the completion of the mission temperatures were again measured at regular intervals during the descent to Riverside. These temperature profiles together with barometric pressure readings obtained on the ground prior to takeoff were later used to reduce the spectrometer strip chart recordings to average pollutant concentrations. The temperature readings were obtained from the aircraft thermometer which has a scale resolution of 1°C.

A summary of the flights and meteorological conditions is presented in Table 1.

3. DATA REDUCTION

The following basic data were used to reduce the spectrometer readings to gas concentration in parts per hundred million by volume:

- (1) Spectrometer readings in the form of continuous graphical recordings of spectrometer output and periodic calibration signals.
- (2) Aircraft ground track from a ground track recovery camera and visual observations.
- (3) Barometric pressure at ground levels from altimeter settings, prior to take off.
- (4) Temperature profiles in the vertical plane.
- (5) Ground elevation along the ground track from topographical maps.
- (6) Sun zenith angles as a function of time over the duration of the mission.
- (7) Visual observations recorded during flight.

Refer to Chart 1 showing the results of Flight 2 for NO₂. The profile shown uses the ground track as a base line and was taken directly from the spectrometer strip chart. It represents the instantaneous total burden (integrated)

concentration pathlength product) of NO_2 as seen by the instrument and is in units of parts per million meters. To reduce ppm-meters to the average concentration in ppm it is necessary to determine the net effective pathlength or thickness of pollutant through which the light has passed before reaching the instrument. The first step in this procedure is to determine the thickness of the gas layer below the aircraft from a knowledge of the elevation of the ground and the measured or observed height of pollutant layer. For the purposes of this program the ground elevations were obtained from topographical maps of the Los Angeles area. To simplify the analysis the data run was divided into zones (shown on Flight 2 as I, II, III, IV and V), and a mean ground elevation determined for each zone. The data applicable to each zone is shown in Table 2. The top of the mixing layer was obtained from the inversion height of 6,500 feet recorded on the temperature surrounding shown on Table 2 and the difference between the inversion height and the average ground elevation taken as the mixing depth.

The concentration may therefore be determined from the relation:

$$\text{concentration (ppm)} = \frac{\text{instrument rdg. (ppm-meters)}}{\text{layer thickness x M factor}} \times \frac{1}{k_{av}}.$$

The M factor takes into account the fact that radiation components reaching the instrument come from all parts of a radiant sky as well as directly from the sun and is fully explained in Appendix I. The $\frac{1}{k_{av}}$ factor takes into account the variations of atmospheric temperature and pressure in the pollutant layer and is fully explained in Appendix II. Referring again to Table 2, the data listed as INSTR. RDG. is the average reading obtained by graphically integrating the profile in each zone. The data listed as SUN ANGLE are the solar zenith angles corresponding the mean time of each zone* and were used to obtain the M FACTOR from Appendix I. APP GAS is the apparent total gas burden in a vertical column below the aircraft. This is then modified by the $\frac{1}{k_{av}}$ function to arrive at the actual total gas burden in ppm-meters. Dividing through by the mixing depth in meters results in the concentration in ppm. Since only

* Altitude and Azimuth of the Sun for Latitude +34-00, Longitude +118-00, Nautical Almanac Office, U.S. Naval Observatory, Washington, D.C.

mean values of all variables in each zone are used in the reduction, it is useful to consider the profile as a simple concentration profile with a scale equivalent to the average concentration for each zone divided by the mean height of the profile. Concentration in pphm may then be read at any point on the profile using the scale of pphm/cm. This procedure of reduction is identical for all zones and missions discussed herein.

4. DISCUSSION OF RESULTS

Flight 1/NO₂ Chart I - This mission was terminated before any useful data could be obtained because of very heavy cloud cover over the entire area east of Chino.

Flight 2/NO₂ May 22/68 Chart I - In spite of scattered cloud over parts of the route and heavy cloud cover over the central region of the Los Angeles Basin much useful data were obtained along the northern leg of the pattern. Also shown on Chart I are the meteorological and A.P.C.D. stations showing wind direction and magnitude and ground level pollutant measurements.

It should be borne in mind when comparing airborne results with ground readings that the two types of measurements are vastly different. The A.P.C.D. results are hourly averages obtained at a single point on the ground whereas the airborne result is an instantaneous reading of concentration averaged over the entire pollutant layer. The airborne result can be higher or lower than the ground measurement depending on the relative position of the ground monitor and the primary sources of pollution and the spatial distribution of pollutants in the atmosphere.

In general of course pollutants concentrations can be expected to be higher near the ground and near the sources of emission before they diffuse into the lower levels of the atmosphere or are dispersed by prevailing winds.

In spite of these major differences in the two methods some useful comparisons can be made. Referring again to Chart I, airborne values are shown at selected points along the profile indicated by the symbol Δ . These are points where the ground track and the prevailing wind permit most meaningful comparisons of airborne and ground measurements. The most significant fact in these data is the trend of values from Riverside to the Chatsworth Reservoir. Whereas the A.P.C.D. results increase from 0 to 7 pphm, the airborne results increase from 0.8 pphm to 2.5 pphm. Unfortunately heavy overcast prevented completion of the data run over Los Angeles center and attempts to fly under the overcast resulted in severe turbulence. The substantial quantities of NO_2 measured off the coast near Hawthorne can be attributed to pollution being carried out over the water by the wind. Note also on this Leg 3 of the flight that the western half of the leg is relatively free from pollution since this zone is downwind of rural areas.

Flight 3/ SO_2 Chart II - Heavy cloud formation again limited the survey. Substantial levels of SO_2 were recorded with maximum values in the San Fernando Valley, Los Angeles center and north of Long Beach.

Flight 4/ NO_2 Chart III - Interesting features of this profile are the rapid build up of NO_2 at Long Beach and the increase in pollutant over Riverside between the start of the survey at 16:44 and the end of the mission at 17:38. The increase is probably due to the transport of NO_2 from the Monrovia area to Riverside by the northwest wind.

Flight 5/ SO_2 Chart IV - Surprisingly high levels of SO_2 were recorded with maximum levels obtained downwind of Los Angeles center in the Monterey Park area.

Flight 6/NO₂ Chart V - The excellent repeatability of the NO₂ measurement is demonstrated by three consecutive runs over the north-south leg. Of particular significance is the repeatable rapid build-up of NO₂ inside the Long Beach breakwater and the gradual build-up of NO₂ from the Santa Monica coastline to San Gabriel.

5. COMPARISON OF AIRCRAFT WITH SURFACE DATA

During the course of the flights, the amount of NO_2 as measured in the surface air varied from zero to 10 pphm, as measured by the Los Angeles Air Pollution Control District (L.A. A.P.C.D.). The corresponding concentration of SO_2 was between 1 and 2 pphm.

The times the aircraft overflew various L.A.A.P.C.D. stations are listed in Table 19 together with the relevant NO_2 concentrations measured by the A.P.C.D. In the last column of the table 18 is given the average concentration in the air below the aircraft as measured by the correlation spectrometer.

The results tabulated in the last two columns of Table 18 are plotted out in Figure 1.

A complete analysis of this data is not possible without a full knowledge of the vertical distribution of atmospheric NO_2 . The concentration of pollutant usually falls off with height and therefore we expect the average concentration to be less than the surface concentration. Figure 1 shows how the average concentration was about one third of the surface value.


6. CONCLUSIONS

The airborne survey in the Los Angeles Basin conducted with the Barringer Correlation Spectrometer clearly demonstrates the capability of the equipment as a reliable tool for the synoptic survey of the distribution of total atmospheric pollution of NO_2 or SO_2 and for the measurement of atmospheric diffusion parameters.

During the survey the significantly large variation of NO_2 made it possible to compare average atmospheric pollution with that recorded at the ground truth stations operated by the Los Angeles Air Pollution Control District. The correlation between these two independent methods of measurement was good, bearing in mind normal meteorological variability.

The airborne equipment operated in a reliable manner during the survey and the ferry period to and from Toronto.

PREPARED BY:


A. J. Moffat, Project Engineer

APPROVED BY:


J. MacDowall, General Manager, Research
and Development Division

AJM:jmb

September 24, 1968

SUMMARY OF FLIGHTS

TABLE 1

<u>DATE</u>	<u>FLT. NO.</u>	<u>GAS</u>	<u>LINE NO.</u>	<u>TIME UP P.S.T.</u>	<u>TIME DOWN P.S.T.</u>	<u>REMARKS</u>
21/5	1	NO ₂	1	11:45	13:10	To Chino Airfield, Flt. aborted. Heavy Cu. coverage whole area. Return Riv. Calibration over Riv.
22/5	2	NO ₂	1	07:30	9:34	To Chatsworth Reservoir Broken Cu. enroute.
	2		2			To Santa Monica Coast
	2		3			To South Side L.A. Airport. Sewage Pipe No. II.
	2		4			Sewage Pipe No. II to Hawthorne to Downy N.D.B. Divert to Long Beach Bro- ken to overcast heavy Cu. tops at 5500 feet.
	2		6			Descended to 2500 ft. Leg aborted due to heavy Cu. at tops at 9500 ft. Unable to maintain VFR conditions
23/5	3	SO ₂	1			Broken Cu. conditions
	3		2	11:02	13:37	Clear
	3		3			Clear
	3		4			Scattered to broken Cu.
	3		5			Scattered to broken Cu.
	3		6			Scattered to broken Cu. Tops at 6500 ft.

TABLE 1 (cont'd)

<u>DATE</u>	<u>FLT. NO.</u>	<u>GAS</u>	<u>LINE NO.</u>	<u>TIME UP P.S.T.</u>	<u>TIME DOWN P.S.T.</u>	<u>REMARKS</u>
23/5	4	NO ₂	4&5 comb.	15:30	17:00	To Chino-Pimona Vortac. Radial 200° heading 220° to Long Beach Harbor. Broken to overcast Cu. Over other areas - also over Santa Ana.
	4		6			Long Beach to Anaheim.
24/5	5	SO ₂	7	10:25	13:30	(3) Long Beach to Griffith Park (2) Griffith Park to Long Beach Scattered Cu.
	5		8			(2) Santa Monica to Monterey Park area (1) Monterey Park area to Santa Monica Broken Cu. at Monterey
24/5	6	NO ₂	7	15:30	18:00	(2) Long Beach to Griffith Park (1) Griffith Park to Long Beach
	6		8			(1) Santa Monica to Monterey Park area (1) Monterey Park area to Santa Monica Very poor visibility due to flying into sun

DATA REDUCTION

LOCATION		Los Angeles				
DATE		May 22, 1968				
FLIGHT NUMBER		2				
INVERSION HEIGHT		6,500 Feet				
POLLUTANT		NO ₂				
ZONE		I	II	III	IV	V
TIME P.S.T.	START-STOP	7.50-7.54	7.59-8.06	8.06-8.16	8.20-8.25	8.32-8.37
GROUND ELEVATION	FEET	710	750	775	1100	0
MIXING DEPTH	FEET	5790	5750	5725	5400	6500
INSTR. RDG.	PPM-M	40	59	69	39	19
SUN ANGLE	DEGREES	54	52	50	48	45
M FACTOR		2.76	2.70	2.65	2.61	2.55
APP. GAS	PPM-M	14.5	21.8	26.0	14.9	7.45
1/K _{av} .		1.06	1.06	1.06	1.06	1.06
ACTUAL GAS	PPM-M	15.4	23.1	27.6	15.8	7.9
CONCENTRATION	PPHM	0.87	1.32	1.57	0.96	0.40
SCALE	PPHM/CM.	0.87	0.89	0.91	0.98	0.83

TABLE 2

TABLE 3

TEMPERATURE SOUNDINGS - FLIGHT NO. 2, NO₂

<u>DATE</u>	<u>TIME(P.S.T.)</u>	<u>TEMP. °C</u>	<u>ALTITUDE-FT.</u>	AVERAGE ASC/DESC
				<u>TEMP °C</u>
May 22	07:30	15	790 (on ground)	15.5
		13	1000	13
		12	1500	13
		10	2000	11
		9	3000	9.5
		6	4000	7
		4	5000	5
		4	6000	4
		5	6500	4.5
	07:50	4	6640	
May 22	9:23	6	7000	
		4	6500 - Inversion	
		4	6000	
		5	5500	
		6	5000	
		8	4000 ← Cloud Layer	
		9	3500	
		10	3000	
		11	2500	
		12	2000	
		14	1500	
	9:34	16	Ground, 21° still air	

AND CORRECTION FACTOR CALCULATION

LOCATION: L.A. RIVERSIDE

FLIGHT NO. 2

DATE: May 22

TIME OF TAKE OFF/LAND		ASCENT 07:30 P.S.T.	DESCENT 9:34 P.S.T.
AIRPORT GRD PRESS		in Hg.	29.91"
" " TEMP (on taxi)		°C	15°C
" WIND (GROUND)		mph deg	2 mph @ 260° Magnetic
" " (ALOFT)		knots deg	
(A)	(E)	(F)	(B)
			(C)
			(D)
			(G)
			(H)

[illegible]
$$\Sigma t = 190$$
$$\Sigma kt = 178.9$$
$$K_{AV} = .94$$
$$\frac{1}{K_{av}} = 1.06$$

DATA REDUCTION

LOCATION		Los Angeles					
DATE		May 23, 1968					
FLIGHT NUMBER		3					
INVERSION HEIGHT		5,000 Feet					
POLLUTANT		SO ₂					
ZONE		I	II	III	IV	V	
TIME P.S.T.	START-STOP	1129-1133	1142-1147	1147-1151	1155-1201	1223-1229	
GROUND ELEVATION	FEET	980	1320	875	1100	62	
MIXING DEPTH	FEET	4020	3680	4125	3900	4938	
INSTR. RDG.	PPM-M	61	63	80	40	53	
SUN ANGLE	DEGREES	16	15	14.5	14	16	
M FACTOR		2.47	2.46	2.43	2.50	2.49	
APP. GAS	PPM-M	24.7	25.6	33.0	16.0	21.3	
1/K _{av} .		1.04	1.04	1.04	1.05	1.05	
ACTUAL GAS	PPM-M	25.9	26.6	34.3	16.8	22.4	
CONCENTRATION	PPHM	2.11	2.37	2.73	1.41	1.49	
SCALE	PPHM/CM.	1.39	1.50	1.36	1.41	1.13	

TABLE 5

LOCATION	Los Angeles
DATE	May 23, 1968
FLIGHT NUMBER	3 cont.
INVERSION HEIGHT	5,000 feet
POLLUTANT	SO ₂

NOTE: $\frac{1}{k_{av}}$ from ascent used for Zones II & III

$\frac{1}{k_{av}}$ from average profile used for Zones IV, V & VI

$\frac{1}{k_{av}}$ from descent profile used for Zones VII, VIII & IX

ZONE		VI	VII	VIII	IX	
TIME P.S.T.	START-STOP	1251-1255	1302-1311	1311-1317	1317-1321	
GROUND ELEVATION	FEET	25	35	300	635	
MIXING DEPTH	FEET	4975	4965	4700	4365	
INSTR. RDG.	PPM-M	71	48	30	30	
SUN ANGLE	DEGREES	20	22	24	25	
M FACTOR		2.48	2.53	2.57	2.61	
APP. GAS	PPM-M	28.7	19.0	11.7	11.5	
1/K _{av} .		1.05	1.06	1.06	1.06	
ACTUAL GAS	PPM-M	30.4	20.1	12.4	12.2	
CONCENTRATION	PPM	2.00	1.32	0.86	0.92	
SCALE	PPM/CM.	1.12	1.10	1.15	1.23	

TABLE NO. 6

TEMPERATURE SOUNDING - FLIGHT NO. 3, SO₂

<u>DATE</u>	<u>TIME (P.S.T.)</u>	<u>TEMP. °C</u>	<u>ALTITUDE, FT.</u>	AVERAGE ASC/DESC
				<u>TEMP °C</u>
May 23	11:04	(17)	790 (on ground)	(18.5)
		15	1500	16.5
		13	2000	15
		12	2500	14
		10	3000	12
		9	3500	11.5
		8	4000	9
		6	5000	7 - Inversion
		6	6000	8
	11:14	7-9	7000	8

() = estimated value

May 23	13:27	8	7000
		10	6000
		7	5500
		8	5000
		10	4000
		12	3500
		14	3000
		16	2500
		17	2000
	13:37	18	1500
		(20)	790 (on ground)

TEMP/PRESS PROFILE CHART
AND CORRECTION FACTOR CALCULATION

LOCATION: LOS ANGELES RIVERSIDE

FLIGHT NO. 3, SO₂

DATE. May 23, 1968

[illegible]
$$\begin{aligned}\Sigma t &= 140 \text{ mb} \\ \Sigma kt &= 134.3 \text{ mb}\end{aligned}$$

$$\frac{1}{K_{AV}} = 1.04$$

TEMP/PRESS PROFILE CHART
AND CORRECTION FACTOR CALCULATION

LOCATION: LOS ANGELES RIVERSIDE

FLIGHT NO. 3, SO₂

DATE: May 23, 1968

[illegible]
$$\Sigma t = 140 \text{ mb}$$
$$\Sigma k_t = 133.2 \text{ mb}$$
$$K_{AV} = .95$$
$$\frac{1}{k_{av}} = 1.05$$

TEMP/PRESS PROFILE CHART
AND CORRECTION FACTOR CALCULATION

LOCATION: LOS ANGELES RIVERSIDE

FLIGHT NO. 3, SO₂

DATE: May 23, 1968

[illegible]
$$\begin{aligned}\Sigma t &= 140 \\ \Sigma kt &= 132.7\end{aligned}$$
$$K_{AV} = .946$$
$$\frac{1}{k_{av}} = 1.06$$

DATA REDUCTION

LOCATION		Los Angeles				
DATE		May 23, 1968				
FLIGHT NUMBER		4				
INVERSION HEIGHT		4,800 Feet				
POLLUTANT		NO ₂				
ZONE		I	II	III	IV	V.
TIME P.S.T.	START-STOP	1544-1552	1552-1600	1604-1606	1606-1609	1609-1612
GROUND ELEVATION	FEET	650	800	700	450	275
MIXING DEPTH	FEET	4150	4000	4100	4350	4525
INSTR. RDG.	PPM-M	18	60	65	63	80
SUN ANGLE	DEGREES	54	56	58	58	59
M FACTOR		2.76	2.81	2.88	2.88	2.91
APP. GAS	PPM-M	6.5	21.4	22.6	21.8	27.5
1/K _{av} .		1.05	1.05	1.05	1.05	1.05
ACTUAL GAS	PPM-M	6.8	22.5	23.7	22.9	28.9
CONCENTRATION	PPHM	.54	1.85	1.90	1.73	2.10
SCALE	PPHM/CM.	1.20	1.23	1.17	1.10	1.05

TABLE 10

DATA REDUCTION

LOCATION		Los Angeles					
DATE		May 23, 1968					
FLIGHT NUMBER		4 cont.					
INVERSION HEIGHT		4,800 Feet					
POLLUTANT		NO ₂					
ZONE		VI	VII	VIII	IX	X	
TIME P.S.T.	START-STOP	1612-1618	1626-1634	1634-1636	1636-1639	6139-1642	
GROUND ELEVATION	FEET	75	50	150	340	635	
MIXING DEPTH	FEET	4725	4750	4650	4460	4165	
INSTR. RDG.	PPM-M	45	100	73	59	45	
SUN ANGLE	DEGREES	60	63	64	64.5	65	
M FACTOR		2.96	3.08	3.13	3.13	3.25	
APP. GAS	PPM-M	15.2	32.5	23.4	18.9	13.9	
1/K _{av} .		1.05	1.05	1.05	1.05	1.05	
ACTUAL GAS	PPM-M	16.0	34.1	24.6	19.8	14.6	
CONCENTRATION	PPHM	1.11	2.35	1.74	1.45	1.15	
SCALE	PPHM/CM.	0.99	0.94	0.95	0.98	1.03	

TABLE 10 (CONTINUED)

TABLE 11

TEMPERATURE SOUNDING - FLIGHT NO. 4, NO₂

<u>DATE</u>	<u>TIME (P.S.T.)</u>	<u>TEMP. °C</u>	<u>ALTITUDE, FT.</u>	<u>AVERAGE ASC/DESC TEMP °C</u>
May 23	15:30	22	790 (on ground)	22
		20	1000	19
		18	1500	17.5
		16	2000	16
		14	2500	14
		13	3000	13
		12	3500	12.5
		10.5	4000	10.8
		10	5000	10
		9	6000	9.2
		8	7000	8
		6	8000	
	15:44			
May 23	16:46	8	7500	
		8	7000	
		9	6500	
		9.5	6000	
		10	5500	
		10	5000	
		10	4500	4800 Inversion
		11	4000	
		13	3500	
		13	3000	
		14	2500	
		16	2000	
		17	1500	
		18	1000	
	16:58			

DATA REDUCTION

LOCATION		Los Angeles				
DATE		May 24, 1968				
FLIGHT NUMBER		5				
INVERSION HEIGHT		4000 Feet				
POLLUTANT		SO ₂				
ZONE		I	II	III		
TIME	START-STOP	1103-1105	1105-1111	1234-1243		
GROUND ELEVATION	FEET	400	100	200		
MIXING DEPTH	FEET	3600	3900	3800		
INSTR. RDG.	PPM-M	75	80	100		
SUN ANGLE	DEGREES	18	16.5	16.5		
M FACTOR		2.46	2.44	2.42		
APP. GAS	PPM-M	30.5	32.8	41.3		
1/K _{av} .		1.04	1.04	1.04		
ACTUAL GAS	PPM-M	31.5	36.2	45.6		
CONCENTRATION	PPIM	2.90	3.04	3.94		
SCALE	PPIM/CM.	1.55	1.52	1.58		

TABLE 13

TABLE 14
TEMPERATURE SOUNDING - FLIGHT NO. 5, SO₂

<u>DATE</u>	<u>TIME (P.S.T.)</u>	<u>TEMP. °C</u>	<u>ALTITUDE, FT.</u>	<u>AVERAGE ASC/DESC TEMP °C</u>
May 24	10:23	22	790 (on ground)	22
		18	1000 straight climb to coast	19.5
		16	1500	18
		14	2000	16
		12	2500	14.5
		11	3000	13.5
		10	3500	12
		10	4000 - INVERSION	12
		11	5000	11
		10	6500	10
		10	7000	10
		9	8000	9
	10:32			
May 24	13:20	13	4500	
		14	4000	
		14	3500	
		16	3000	
		17	2500	
		18	2000	
		20	1500	
	13:28	21	1000	

TABLE 15

TEMP/PRESS PROFILE CHART
AND CORRECTION FACTOR CALCULATION

LOCATION: RIVERSIDE MUNICIPAL

FLIGHT NO. 5, SO₂

DATE: May 24, 1968

[illegible]

$\Sigma t = 110$

$$\Sigma k_t = 105.9$$
$$K_{AV} = .962$$
$$\frac{1}{k_{av}} = 1.04$$

DATA REDUCTION

LOCATION		Los Angeles				
DATE		May 24, 1968				
FLIGHT NUMBER		6				
INVERSION HEIGHT		3,000 Feet				
POLLUTANT		NO ₂				
ZONE		I	II	III	IV	
TIME	START-STOP	1633-1647	1558-1610	1614-1625	1703-1710	
GROUND ELEVATION	FEET	160	160	160	200	
MIXING DEPTH	FEET	2840	2840	2840	2800	
INSTR. RDG.	PPM-M	65	80	80	95	
SUN ANGLE	DEGREES	64.5	57	60.5	70	
M FACTOR		3.18	2.84	2.95	3.52	
APP. GAS	PPM-M	20.4	28.2	27.1	27.0	
1/K _{av} .		1.03	1.03	1.03	1.03	
ACTUAL GAS	PPM-M	21.0	29.1	27.9	27.8	
CONCENTRATION	PPIM	2.42	3.36	3.22	3.26	
SCALE	PPIM/CM.	1.49	1.68	1.61	1.37	

TABLE 16

TABLE 17
TEMPERATURE SOUNDING - FLIGHT NO. 6, NO₂

<u>DATE</u>	<u>TIME (P.S.T.)</u>	<u>TEMP. °C</u>	<u>ALTITUDE, FT.</u>	<u>AVERAGE ASC/DESC TEMP. °C</u>
May 24	15:30	26, still	790 (on ground)	
		22	1000	
		20	1500	
		20	2000-3000, heavy air traffic	
		17	3500	
		16	4000	
		15	4500	
		13	5000	
		12	6000	
		10	7500	
		8	8000	
		7	10,000	
	15:58	6	10,300	
	17:40	7	10,000	(7)
		8	9500	(8)
		9	9000	
		10	8500	
		10	8000	9
		12	7000	
		13	6000	12.5
		14	5500	
		15	4500	15
		17	4000	16.5
		18	3500	17.5
		16	3000 - INVERSION	16 - INVERSION
		17	2500	17
		17	2000	17
		18	1500	19
	18:00	18	1000	20
		20	790 (on ground)	23

() = estimated value

TEMP/PRESS PROFILE CHART
AND CORRECTION FACTOR CALCULATION

LOCATION: RIVERSIDE MUNICIPAL

FLIGHT NO. 6, NO₂

DATE: May 24, 1968

[illegible]
$$\begin{aligned}\Sigma t &= 77 \\ \Sigma kt &= 74.7\end{aligned}$$
$$K_{AV} = .97$$
$$\frac{1}{k_{av}} = 1.03$$

TABLE 19 - COMPARISON BETWEEN AIRCRAFT AND GROUND NO₂ DATA

May 22, 1968

<u>TIME (P.S.T.)</u>	<u>STATION</u>	<u>A.P.C.D. POLLUTION AT SURFACE (pphm)</u>	<u>BARRINGER AVERAGE POLLUTION (pphm)</u>
7.52	Chino	0	0.8
7.59	73	2	0.4
8.02	60	1	1.1
8.05	64	4	2.1
8.11	69	6	1.3
8.15	74	7	2.5


May 23, 1968

15.56	75	7	2.5
16.14	72	7	1.5
16.30	72	7	2.3

May 24, 1968

16.42	72	10	3.7
16.04	72	10	3.0
16.20	72	10	3.5
17.07	1	7	4.1

FIGURE 1 - COMPARISON BETWEEN AVERAGE POLLUTION AND SURFACE POLLUTION



APPENDIX A - DERIVATION OF PATHLENGTH FACTOR

1) RADIATION COMPONENTS

For a study concerned with the radiation arriving at an aircraft-placed spectrometer looking directly down, it is necessary to separate the radiation incident on the ground from a source point of view.

First, we consider the direct sun component, defined by its spectral irradiance in $\text{w cm}^{-2} \text{ nm}^{-1}$ for a given wavelength.

It will be denoted as H_λ , and in a general case we expect it to be a function $H_\lambda = H_\lambda(\theta, \tau, T, \psi)$ of the following parameters

θ	sun zenith angle
ψ	air mass
τ	ozone thickness
T	atmospheric turbidity

The spectral radiant power on the ground unit area P_λ in w nm^{-1} will be

$$P'_\lambda = H_\lambda \cos\theta \quad (1-1)$$

Considering a Lambertian reflection from the ground of reflecting $R_f(\lambda)$, the spectral radiance of the ground as observed from the aircraft-based instrument

$$\text{is} \quad N'_\lambda = \frac{R_f(\lambda) H_\lambda \cos\theta}{\pi} \quad \text{w cm}^{-2} \text{ sr}^{-1} \text{ nm}^{-1} \quad (1-2)$$

The second component, that of the radiation emerging from the atmosphere due to the scattering of the sunlight is

$$N'_\lambda = N'_\lambda [\theta, \tau, \psi, \phi, \gamma, R_f(\lambda)]$$

for a perfect Rayleigh atmosphere, where:

θ is the sun zenith angle
 τ ozone thickness
 ψ air mass
 ϕ azimuth of emission area referred to a vertical plane passing through the local zenith and the sun zenith angle of emission area
 γ area
 γ zenith angle on emission area

$R_f(\lambda)$ reflectivity of ground

For a real atmosphere and a given specific location and time, this component is more likely to be

$$N'_\lambda = N'_\lambda (\theta, \tau, \psi, \phi, \gamma, R_f(\lambda) T)$$

where T is the atmospheric turbidity, function of the aerosol content, size and distribution.

The spectral radiant power on the ground due to a differential area of the sky that is projected into the solid angle $\frac{1}{R_2} \cos \gamma$ [1 = unit area on the ground-cm²
R = imaginary distance from sky area to the ground] by which the unit ground area is seen from the sky area, is:

$$P''_\lambda = N'_\lambda \cos \gamma \sin \gamma \, d\gamma \, d\phi \quad \text{w nm}^{-1} \quad (1-3)$$

The spectral radiance of the ground unit area as observed from the aircraft and from this second component is

$$N''_\lambda = \frac{R_f(\lambda)}{\pi} \iint N'_\lambda \cos \gamma \sin \gamma \, d\gamma \, d\phi \quad \text{w cm}^{-2} \text{ sr}^{-1} \text{ nm}^{-1} \quad (1-4)$$

for a fixed λ

The total energy arriving at the instrument is

$$P_{\lambda} = A\Omega \left[\int_{\lambda_1}^{\lambda'_1} N_{\lambda} d\lambda + \int_{\lambda_1}^{\lambda'_1} N'_{\lambda} d\lambda \right] \quad \text{in watts} \quad (1-5)$$

A = instrument aperture

Ω = instrument acceptance angle

$\lambda_1 - \lambda'_1$ = waveband of interest

where no absorption or scattering below the aircraft are considered.

2) CORRELATION SPECTROMETRY APPLICATION

In this section we are concerned with the absorption of the gas under study.

Our assumptions are:

- 1st The absorption is done only by the gas
- 2nd There is no scattering below the aircraft
- 3rd All the gas is below the aircraft
- 4th The instrument optics are such that the response to all wavelengths within the interval of interest is linear-constant

For the general case of an airborne-based instrument looking vertically down, the center of the ground patch determined by its field of view is taken as the origin of coordinates.

The vertical axis is that of the local zenith, the azimuth angle (ϕ) is measured with reference to a plane passing through the vertical and the sun, and the angle γ is measured from the zenith.

The concentration of the absorbing gas for the line of sight $S(\phi, \gamma)$ is in the most general case a function $c = c(\phi, \gamma, \ell)$ (for a given period of time) of those angles and another independent variable (ℓ). Here the length along the line of sight is taken as such.

The beam of light proceeding from the fictitious zone in the sky of area $R^2 \sin \gamma d\gamma d\phi$ and spectral radiance N'_λ travels along the pathlength ℓ and is attenuated by

$$e^{-a(\lambda) \cdot \int_0^\ell c d\ell} \quad (\text{for every wavelength})$$

$a(\lambda)$ being the absorption crosssection per molecule, function of wavelength

After a Lambertian reflection on the ground of reflectivity $R_f(\lambda)$, it is attenuated by

$$e^{-a(\lambda)} \int_0^h c_{\lambda} dz$$

on its way back to the aircraft.

c_{λ} corresponding to $c(0, 0, h)$ depends only on height (h) .

The elemental energy in the instrument due to the beam travelling this path is:

$$dP = \frac{A \Omega}{\pi} \cdot R_f(\lambda) N_{\lambda}' e^{-a(\lambda) \left[\int_0^{\ell} c d\ell + \int_0^h c_1 dz \right]} \sin \gamma \cos \gamma d\gamma d\varphi d\lambda \quad (2-1)$$

The energy in the instrument proceeding from the whole sky for one slit is

$$P = \frac{A \Omega}{\pi} \iiint R_f(\lambda) N_{\lambda}' e^{-a(\lambda) \left[\int_0^{\ell} c d\ell + \int_0^h c_1 dz \right]} \cos \gamma \sin \gamma d\gamma d\varphi d\lambda \quad (2-2)$$

λ integrated for the limits defining the slit

The other component, that due to the direct sunlight following the same process, will be

$$dP_2 = \frac{A \Omega}{\pi} R_f(\lambda) H_{\lambda} \cos \theta e^{-a(\lambda) \left[\int_0^{\ell} c_2 d\ell + \int_0^h c_1 dz \right]} d\lambda$$

$$P_2 = \frac{A \Omega}{\pi} \int R_f(\lambda) H_{\lambda} \cos \theta e^{-a(\lambda) \left[\int_0^{\ell} c_2 d\ell + \int_0^h c_1 dz \right]} d\lambda \quad (2-3)$$

C_2 is the concentration along the line of sight $S(0, \theta)$ corresponding to $c(0, \theta, \lambda)$

The modulation at an aircraft-based instrument looking down will be

$$M = \frac{A\Omega}{\pi} \left\{ \sum_{i=1}^m \left[\iiint R_f(\lambda) N'_\lambda \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] \cos \gamma \sin \gamma d\gamma d\varphi d\lambda + \right. \right. \\ \left. \left. \int R_f(\lambda) H_\lambda \cos \theta \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] d\lambda \right]_{\lambda_{1i}}^{\lambda_{2i}} - \right. \\ \left. \sum_{i=1}^m \left[\iiint R_f(\lambda) N'_\lambda \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] \cos \gamma \sin \gamma d\gamma d\varphi d\lambda + \right. \right. \\ \left. \left. \int R_f(\lambda) H_\lambda \cos \theta \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] d\lambda \right]_{\lambda_{2i}}^{\lambda_{2i}'} \right\} \frac{\sigma}{\omega}$$

$$\left| \sum_{i=1}^m \left[\iiint R_f(\lambda) N'_\lambda \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] \cos \gamma \sin \gamma d\gamma d\varphi d\lambda + \right. \right. \\ \left. \left. \int R_f(\lambda) H_\lambda \cos \theta \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] d\lambda \right]_{\lambda_{1i}}^{\lambda_{2i}'} \right| \quad (2-4)$$

$$\text{or } M = \frac{\sum_{i=1}^m [P + P_2]_{\lambda_{1i}}^{\lambda_{2i}'} - \sum_{i=1}^m [P + P_2]_{\lambda_{2i}}^{\lambda_{2i}'}}{\left| \sum_{i=1}^m [P + P_2]_{\lambda_{1i}}^{\lambda_{2i}'} \right|} \quad (2-5)$$

$\lambda_{1i} - \lambda_{2i}'$ are the beginning and end wavelength of the mask corresponding to the slit (i) for the position (1) of the mask with respect to the spectrum.

$\lambda_{1i} - \lambda_{2i}$ The same for the next position (2) of the mask with respect to the spectrum.

n The number of slits in the mask.

The purpose of aircraft measurements is to obtain "as accurate as possible" measurement of c_1 , that is, the concentration directly below the aircraft; this concentration cannot be calculated for the more general case.

In aircraft use, the zeroing is done against the vertical and the measurement, looking down. The reading R will be equal to

$$R = \alpha \left\{ \frac{\sum_{i=1}^n \int_{\lambda_{2i}}^{\lambda'_{2i}} (N'_\lambda)_v d\lambda}{\sum_{i=1}^n \int_{\lambda_{1i}}^{\lambda'_{1i}} (N'_\lambda)_v d\lambda} \right\} \quad \text{I}$$

$$\sum_{i=1}^n \left[\iiint R_f(\lambda) N'_\lambda \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] \sin \gamma \cos \gamma d\gamma d\varphi d\lambda + \right. \\ \left. \int R_f(\lambda) H_\lambda \cos \theta \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] d\lambda \right]_{\lambda_{2i}}^{\lambda'_{2i}} \div$$

$$\sum_{i=1}^n \left[\iiint R_f(\lambda) N'_\lambda \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] \sin \gamma \cos \gamma d\gamma d\varphi d\lambda + \right.$$

$$+ \left\{ \int_{\lambda_{1i}}^{\lambda_{2i}} R_f(\lambda) H_\lambda \cos \theta \exp \left[-a(\lambda) \left(\int_0^l c_2 dl + \int_0^h c_1 dz \right) \right] d\lambda \right\}$$

II

(2-6)

$(N'_\lambda)_v$ = spectral radiance of the vertical zone of the sky

α The instrument constant, combination of optics and electronics, transmission and response factors.

The instrument calibration is done against the vertical; therefore, the calibration values follow the expression

$$CR = \alpha \left\{ \frac{\sum_{i=1}^n \int_{\lambda_{2i}}^{\lambda'_{2i}} (N'_\lambda)_v d\lambda}{\sum_{i=1}^n \int_{\lambda_{1i}}^{\lambda'_{1i}} (N'_\lambda)_v d\lambda} - \frac{\sum_{i=1}^n \int_{\lambda_{2i}}^{\lambda'_{2i}} (N'_\lambda)_v e^{-a(\lambda)cl} d\lambda}{\sum_{i=1}^n \int_{\lambda_{1i}}^{\lambda'_{1i}} (N'_\lambda)_v e^{-a(\lambda)cl} d\lambda} \right\} \quad (2-7)$$

III

IV

and we are left with the equality

$$[\text{expression (II)} = \text{expression (IV)}] \quad (2-8)$$

The assumptions that we must make are that

- 1) N'_λ is isotropic; that is, it does not depend on (ϕ, γ) . Then, $N'_\lambda = (N'_\lambda)_{\text{v}}$
- 2) N'_λ can be expressed as a ratio of H_λ i.e. $N'_\lambda = f(H_\lambda)$
and that this function is known, for a given period of time, ozone thickness etc. Then this ratio will be a function of λ and θ only; $N'_\lambda = H_\lambda f(\lambda, \theta)$
- 3) The gas is stratified; that is, its concentration is constant over horizontal layers. And the total concentration x pathlength can be expressed $ch = c_1 h_1 + c_2 h_2 + \dots \dots \dots c_n h_n$
($c_i h_i$ is the concentration on layer (i) times the height of the layer (h_i)
where c_i is constant)
- 4) $Rf(\lambda)$ is constant = R over the interval $\lambda_{11} \leftrightarrow \lambda'_{2n}$

Because of (3) the expression

$$e^{-a(\lambda) \left[\int_0^\ell c \, d\ell + \int_0^h c_1 \, dz \right]}$$

reduces to

$$e^{-a(\lambda) ch \left[\sec \gamma + 1 \right]}$$

and $e^{-a(\lambda) \left[\int_0^\ell c_2 \, d\ell + \int_0^h c_1 \, dz \right]}$ to $e^{-a(\lambda) ch \left[\sec \theta + 1 \right]}$

and expression (2-8) becomes

$$\frac{\sum_{i=1}^n \int_{\lambda_{2i}}^{\lambda'_{2i}} H_{\lambda} F(\theta, \lambda) e^{-a(\lambda)[c\ell]} d\lambda}{\sum_{i=1}^n \int_{\lambda_{1i}}^{\lambda'_{1i}} H_{\lambda} F(\theta, \lambda) e^{-a(\lambda)[c\ell]} d\lambda} =$$

$$\frac{\sum_{i=1}^n \left[\iiint H_{\lambda} F(\theta, \lambda) \exp \left[-a(\lambda) \operatorname{ch}(\sec \gamma + 1) \right] \sin \gamma \cos \gamma d\gamma d\phi d\lambda + \int H_{\lambda} \cos \theta \exp \left[-a(\lambda) \operatorname{ch}(\sec \theta + 1) \right] d\lambda \right]_{\lambda_{2i}}^{\lambda'_{2i}}}{\sum_{i=1}^n \left[\iiint H_{\lambda} F(\theta, \lambda) \exp \left[a(\lambda) \operatorname{ch}(\sec \gamma + 1) \right] \sin \gamma \cos \gamma d\gamma d\phi d\lambda + \int H_{\lambda} \cos \theta \exp \left[-a(\lambda) \operatorname{ch}(\sec \theta + 1) \right] d\lambda \right]_{\lambda_{1i}}^{\lambda'_{1i}}} \quad (2-9)$$

The second term can be further simplified if $f(\lambda, \theta)$ can be expressed as a product of two other functions, each one depending on only one variable, that is $f(\lambda, \theta) = Z(\theta) \cdot L(\lambda)$. The limits of integration of ϕ and γ are:

$$\phi \quad 0 \leftrightarrow 2\pi$$

$$\gamma \quad 0 \leftrightarrow \frac{\pi}{2}$$

and the expression (2-8) becomes

$$\frac{\sum_{i=1}^m Z(\theta) \int_{\lambda_{2i}}^{\lambda'_{2i}} H_{\lambda} L(\lambda) e^{-a(\lambda)[cl]} d\lambda}{\sum_{i=1}^m Z(\theta) \int_{\lambda_{1i}}^{\lambda'_{1i}} H_{\lambda} L(\lambda) e^{-a(\lambda)[cl]} d\lambda} = \frac{\sum_{i=1}^m \int_{\lambda_{2i}}^{\lambda'_{2i}} H_{\lambda} L(\lambda) e^{-a(\lambda)[cl]} d\lambda}{\sum_{i=1}^m \int_{\lambda_{1i}}^{\lambda'_{1i}} H_{\lambda} L(\lambda) e^{-a(\lambda)[cl]} d\lambda} =$$

(A) (B)

$$\frac{\sum_{i=1}^m \left[2\pi Z(\theta) \iint H_{\lambda} L(\lambda) \exp[-a(\lambda)ch(\sec\theta+1)] \sin\theta \cos\theta d\gamma d\lambda + \int H_{\lambda} \cos\theta \exp[-a(\lambda)ch(\sec\theta+1)] d\lambda \right]_{\lambda_{2i}}^{\lambda'_{2i}}}{\sum_{i=1}^n \left[2\pi Z(\theta) \iint H_{\lambda} L(\lambda) \exp[-a(\lambda)ch(\sec\theta+1)] \sin\theta \cos\theta d\gamma d\lambda + \int H_{\lambda} \cos\theta \exp[-a(\lambda)ch(\sec\theta+1)] d\lambda \right]_{\lambda_{1i}}^{\lambda'_{1i}}}$$

(C)

(2-10)

We name the value $CR = \alpha(1-B)$ [expression (2-7) after having been simplified with the given assumptions], the calibrated reading corresponding to the concentration $[cl]$.

The problem can be stated as follows:

Which concentration ch below the aircraft would, after following the studied process, give a reading in the calibration curve corresponding to $[cl]$?

The way to solve it is as follows: Knowing $Z(\theta)$, $L(\lambda)$, H_{λ} and $a(\lambda)$, introduce values of ch and $[cl]$ in sides B and C of equation (2-10), and see which value of ch gives a value to the term C equal to a $[cl]$ in the term B.

The values ch , $[cl]$ in ppm-M can be then entered in a table or tables, for each value of the angle θ . In that situation $[cl]$ would be the "readout calibrated concentration" for the angle θ due to the real concentration ch below the aircraft. Then, the concentration ch could be determined with a set of tables for different times of year, geographical locations and sun angles.

If we further assume that

- (a) $H\lambda$ has a continuous distribution over the waveband of interest.
- (b) $L(\lambda)$ is equal to 1, that is the spectral distribution of $H\lambda$ is equal to that of $N\lambda$.
- (c) The width of all slits in the mask is the same.
- (d) Two values a_1 and a_2 of $a(\lambda)$ can be found, such that expression (2-10) becomes

$$e^{-(a_2 - a_1)[c\lambda]} = \frac{e^{-a_2 \operatorname{ch}(\sec\theta + 1)} + 2\pi Z(\theta) \int_0^{\frac{\pi}{2}} e^{-a_2 \operatorname{ch}[\sec\gamma + 1]} \cos\gamma \sin\gamma d\gamma}{e^{-a_1 \operatorname{ch}(\sec\theta + 1)} + 2\pi Z(\theta) \int_0^{\frac{\pi}{2}} e^{-a_2 \operatorname{ch}[\sec\gamma + 1]} \cos\gamma \sin\gamma d\gamma}$$

(2-11)

For the solution of this formula, the value of $\pi \cdot Z(\theta)$

has been taken as $-\frac{450}{8(\theta - 76.25)}$ for SO_2

and as $-\frac{42}{\theta - 140}$ for NO_2 .

θ , in degrees,

as it was explained in TR-68-56 , "Final Report, Absorption Spectroscopy Experiment", work done on NASA Contract No. NAS9-7241.

A computer program (PATH) has been written to yield values of

$$m = \frac{[c\ell]}{ch}$$

using the values,

$$\text{NO}_2 \quad \left\{ \begin{array}{l} a_2 = .332 * 10^{-18} \frac{\text{cm}^2}{\text{molec.}} \\ a_1 = .166 * 10^{-18} \frac{\text{cm}^2}{\text{molec.}} \end{array} \right.$$

and


$$\text{SO}_2 \quad \left\{ \begin{array}{l} .73 * 10^{-18} \frac{\text{cm}^2}{\text{molec.}} \\ .34 * 10^{-18} \frac{\text{cm}^2}{\text{molec.}} \end{array} \right.$$

Prepared by



M. Millan
Mathematician

Approved by



J. MacDowall
General Manager
Research and Development Division

September 30, 1968

APPENDIX B

AIRBORNE SO₂/NO₂ MEASUREMENTS CALCULATION OF CONCENTRATION FROM TOTAL VERTICAL GAS BURDEN

The total vertical gas burden is considered to be that component of the spectrometer read-out which represents the concentration X pathlength product of the pollutant integrated along a vertical line of sight from ground level to the top of the pollutant layer. Given this quantity plus the temperature at various points throughout the layer and barometric pressure at ground level, the problem is to find the average concentration in the layer.

It should be borne in mind in this discussion that the spectrometer readout is quantitized in flight by calibrations of instrument sensitivity using a reference cell of known gas burden. The spectrometer contains two reference cells (for two different gases) which have been carefully measured or calibrated in the laboratory prior to the survey. For this particular mission the NO₂ cell contained 85.8 ppm-meters and the SO₂ cell 155 ppm-meters of gas-both at 20°C and one atmosphere of pressure.

The integrated vertical burden c_l in ppm-meters has concentration c in parts per million by volume (i.e., by no. of molecules) at 1 atmosphere and 20°C (293°A). That is, that number of molecules per unit volume of a gas at 20°C and 1 atmosphere would have c parts per million. Now consider the density of air at 20°C and 1 atmosphere to be 1. At a different temperature T and pressure P the air will have a different density $d = \frac{T_0}{T} \times \frac{P}{P_0} = \frac{293^\circ A}{T} \times \frac{P}{1 \text{ atmos.}}$. If the density is lower (higher) the given amount of pollutant will form a higher (lower) fraction in parts per million of the total.

We assume that the pollutant layer is homogenous, i.e., the pollutant forms a fraction of the layer which is constant with height. The density of the pollutant varies in proportion to the overall density of the air. We divide

the pollutant layer into thin horizontal sub-layers or slices. We assume a concentration of 1 for each slice in parts per million at the temperature and pressure of the slice. We then find the concentration reduced to 20°C and 1 atmosphere. We then find the average concentration k (at 20°C + 1 atm.) of the slices weighted according to the amount of air in each slice.

Now we find the height z between the ground and the top of the pollutant layer. If the integrated burden or thickness is kz ppm-m, then the average concentration reduced to 20°C and 1 atm. is k . Also if cz is the actual integrated thickness, that is, the total vertical burden in ppm-meters at ambient temperature and pressure, then $c \times \frac{1}{k}$ is the concentration. Now will be described a method to find $\frac{1}{k}$

Refer now to the pollution survey conducted by Barringer Research in Los Angeles on May 22. Refer further to the temperature sounding taken on the ascent. For the purpose of this example we will assume the pollutant layer extends from a ground elevation of 600 feet (m.s.l.) to a maximum height of 5000 feet (m.s.l.).

(ground)							Calculate To-	Line	
Height (m.s.l)	600	1000	1500	2000	3000	4000	5000	nearest 100 ft.	A
Pressure	990	980	960	940	905	875	845	nearest 5 mb	B
Slice Thickness t	10	20	20	35	30	30		nearest 5 mb	C
Rel. Pressure $\frac{P}{1 \text{ atm.}}$.97	.955	.935	.91	.88	.85		nearest .005	D
Temp. T	287	285.5	284	282.5	280.5	278		nearest 5°A	E
$\frac{T}{293^{\circ}\text{A}}$.98	.975	.97	.965	.96	.95		nearest .005	F
$\frac{P}{1 \text{ atm.}} \times \frac{293^{\circ}\text{A}}{T} = k$.99	.98	.965	.95	.915	.895		nearest .005	G
kt	9.9	19.6	19.3	33.2	27.5	26.8		nearest .1 mb	H
$\Sigma t = 145 \text{ mb}$		$\Sigma kt = 136.3 \text{ mb}$							I
$k_{\text{av.}} = \frac{136.3}{145} = .94 \frac{1}{k_{\text{av.}}} = \frac{145}{136.3} = 1.065$									J

- Line A - height above mean sea level recorded by altimeter at each point at which reading is taken from bottom to top of pollution layer. Readings should be taken at ground and every 1000 feet thereafter until the top of pollution layer. A slice is taken between each pair of levels.
- Line B - pressure in mb. at each recording height. This is taken from a U.S. upper air chart based on the U.S. standard atmosphere. If an altimeter corrected for temperature, or a radar altimeter, is used, a different atmosphere will have to be used for hot or cold weather. Remember to add or subtract a correction factor (to nearest 5 mb) if the sea level pressure is above or below the standard 1013 mb. value. In the example the correction factor is 0.
- Line C - slice thickness t . This is the difference in pressures (from line B) at adjacent recording heights. This gives the amount of air in each slice.
- Line D - relative pressure. Take the mean of the pressures at the endpoints of the slice and divide by the standard mean sea level pressure to get slice pressure in atmospheres. The first value in the example is $(\frac{990 + 980}{2}) \times \frac{1}{1013} = .97$.
- Line E - Temperature T . Take the mean of the temperatures at the endpoints of the slice and convert to absolute. The first value in the example is $\frac{15 + 13}{2} + 273 = 287^{\circ}\text{A}$.
- Line F - relative temperature $\frac{T}{293^{\circ}\text{A}}$
- Line G - factor k for each slice. Divide the value from line D by that from line F.
- Line H - kt . Multiply line C by line G to obtain this intermediate step in finding the average of k .

Line I - Σt (diff. in press. between top and bottom) from line B on (which should give the same answer). Also we have Σkt from line H.

Line J - $k_{av.} = \frac{\Sigma kt}{\Sigma t}$ from line I, $\frac{1}{k_{av.}}$ is also determined.

Suppose the integrated thickness was 120 ppm-m. Divide by the layer thickness 5000-600 = 4400 feet - 1420 meters to get $\frac{120}{1420}$ ppm (20°C, 1 atm.) average volume concentration. Multiply by $\frac{1}{k_{av.}}$ to find the actual average concentration:

$$\frac{120 \times 1.065}{1420} = .090 \text{ ppm}$$

Prepared by



A. J. Moffat
Project Engineer

and

S. Clodman
Scientist

Approved by



Joseph MacDowall
General Manager
Research and Development Division

APPENDIX C
GROUND TRUTH AIR POLLUTION DATA
ACQUIRED BY LOS ANGELES AIR POLLUTION
CONTROL DISTRICT (LAAPCD), LOS ANGELES COUNTY


During the course of the Barringer survey, the concentration of NO_2 and SO_2 in the air at ground level was measured by the LAAPCD. These stations have been in routine operation for 10 to 15 years. The amount of NO_2 in the surface air is measured by the Saltzman reagent method.

The attached figure gives the location of the ground stations. The recorded concentration is given in four tables.

AIR POLLUTION CONTROL DISTRICT
LOS ANGELES COUNTY

SMOG FORECAST AREAS

NOVEMBER 1966



ATMOSPHERIC CONCENTRATIONS OF NO₂ AND SO₂ IN THE LOS ANGELES BASIN

May 21, 1968

No.	STATION Location	Contam- inant	HOURLY AVERAGE CONCENTRATION (PPHM)										6 PM *
			8 AM	9 AM	10 AM	11 AM	12 N	1 PM	2 PM	3 PM	4 PM	5 PM	
1	Central Los Angeles	NO ₂	7	7	6	-	-	5	3	5	5	3	3
		SO ₂	1	1	-	1	1	1	1	1	1	1	1
60	East San Gabriel Valley	NO ₂	3	1	1	1	-	-	-	1	1	1	1
64	West San Gabriel Valley	NO ₂	2	3	3	-	-	-	6	5	3	3	3
		SO ₂	1	1	1	1	1	-	2	2	2	2	2
69	East San Fernando Valley	NO ₂	4	6	7	7	-	-	-	6	6	6	5
		SO ₂	1	1	1	-	-	1	2	1	1	1	1
71	West Los Angeles	NO ₂	3	3	3	-	-	-	3	3	3	3	3
		SO ₂	2	1	-	1	2	2	2	2	2	2	2
72	South Coastal	NO ₂	-	-	-	-	-	-	-	-	-	-	6
		SO ₂	3	3	6	2	2	2	-	1	1	1	1
74	West San Fernando Valley	NO ₂	2	2	2	2	-	-	-	2	2	2	2
		SO ₂	2	1	1	-	1	1	1	1	1	1	1
75	Pomona-Walnut Valley	NO ₂	2	3	3	3	-	-	-	4	5	6	6
		SO ₂	1	1	1	-	1	1	1	1	1	1	2
76	South West Coastal	NO ₂	-	-	4	6	-	-	-	4	5	5	4
		SO ₂	1	1	2	2	2	1	-	2	3	3	2

* All times used are Pacific Standard Time.

ATMOSPHERIC CONCENTRATION OF NO₂ AND CO₂ IN THE LOS ANGELES BASIN

May 22, 1968

No.	STATION Location	Contam- inant	HOURLY AVERAGE CONCENTRATION (PPHM)										
			8 AM	9 AM	10 AM	11 AM	12 N	1 PM	2 PM	3 PM	4 PM	5 PM	6 PM
1	Central Los Angeles	NO ₂	9	9	8	-	5	4	4	4	4	3	3
		SO ₂	3	3	3	2	2	-	2	1	1	1	1
60	East San Gabriel Valley	NO ₂	3	1	1	1	-	-	-	1	1	1	1
64	West San Gabriel Valley	NO ₂	3	4	4	-	-	4	4	3	2	2	2
		SO ₂	2	-	2	-	1	1	1	1	1	1	1
69	East San Fernando Valley	NO ₂	6	6	7	8	-	-	-	4	4	4	7
		SO ₂	-	2	2	2	2	2	2	2	2	2	2
71	West Los Angeles	NO ₂	11	9	-	-	-	4	4	3	3	3	3
		SO ₂	-	2	1	1	1	1	1	1	1	1	1
72	South Coastal	NO ₂	6	5	-	-	-	-	-	-	-	-	-
		SO ₂	2	2	2	2	2	2	2	2	2	2	2
74	West San Fernando Valley	NO ₂	8	7	5	6	-	-	1	1	1	1	1
		SO ₂	2	2	1	1	2	2	2	1	1	1	1
75	Pomona-Walnut Valley	NO ₂	2	2	3	4	-	-	2	3	5	4	3
		SO ₂	1	-	1	1	1	-	1	2	2	2	2
76	South West Coastal	NO ₂	8	8	4	3	5	-	-	3	3	4	3
		SO ₂	3	7	-	2	2	2	1	1	1	1	1

* All times used are Pacific Standard Time.

ATMOSPHERIC CONCENTRATION OF NO₂ AND SO₂ IN THE LOS ANGELES BASIN

May 23, 1968

No.	STATION Location	Contam- inant	HOURLY* AVERAGE CONCENTRATION (PPHM)										
			8 AM	9 AM	10 AM	11 AM	12 N	1 PM	2 PM	3 PM	4 PM	5 PM	6 PM
1	Central Los Angeles	NO ₂	5	6	5	-	5	5	4	5	5	5	4
		SO ₂	2	-	2	2	2	2	2	2	2	2	2
60	East San Gabriel Valley	NO ₂	1	1	1	1	-	-	-	1	1	1	1
64	West San Gabriel Valley	NO ₂	3	2	3	-	-	5	4	3	3	3	3
		SO ₂	1	1	1	-	-	2	2	2	2	2	2
69	East San Fernando Valley	NO ₂	3	3	5	6	-	-	-	4	5	6	6
		SO ₂	2	2	2	2	2	2	2	2	2	2	2
71	West Los Angeles	NO ₂	6	5	-	-	-	3	3	3	4	3	3
		SO ₂	2	2	3	2	2	1	2	2	1	1	1
72	South Coastal	NO ₂	6	5	5	5	-	3	6	7	7	7	5
		SO ₂	1	7	1	1	2	3	2	2	1	1	1
74	West San Fernando Valley	NO ₂	-	-	3	5	4	3	2	2	2	2	2
		SO ₂	1	1	1	1	1	1	1	1	1	1	1
75	Pomona-Walnut Valley	NO ₂	3	2	3	3	-	-	-	8	7	7	7
		SO ₂	1	-	1	1	1	1	1	1	2	2	2
76	South West Coastal	NO ₂	6	6	6	3	-	-	-	6	8	4	2
		SO ₂	1	-	3	2	3	2	1	1	1	1	1

* All times used are Pacific Standard Time.

ATMOSPHERIC CONCENTRATION OF NO₂ AND SO₂ IN THE LOS ANGELES BASIN

May 24, 1968

No.	STATION Location	Contam- inant	HOURLY AVERAGE CONCENTRATION (PPHM)										
			8 AM	9 AM	10 AM	11 AM	12 N	1 PM	2 PM	3 PM	4 PM	5 PM	6 PM*
1	Central Los Angeles	NO ₂	7	9	7	-	-	6	6	7	7	7	7
		SO ₂	3	3	3	2	2	2	2	1	2	2	1
60	East San Gabriel Valley	NO ₂	1	1	1	1	-	-	-	1	1	1	1
64	West San Gabriel Valley	NO ₂	5	4	-	-	-	-	5	4	4	5	6
		SO ₂	2	2	1	1	1	1	1	1	1	1	1
69	East San Fernando Valley	NO ₂	7	7	11	9	-	-	-	6	8	11	14
		SO ₂	2	-	2	2	3	3	2	2	2	2	2
71	West Los Angeles	NO ₂	11	9	-	-	-	3	3	4	4	4	4
		SO ₂	2	2	1	1	1	1	1	1	1	1	1
72	South Coastal	NO ₂	3	3	7	4	-	-	-	3	5	7	10
		SO ₂	1	-	-	1	1	1	1	1	1	1	1
74	West San Fernando Valley	NO ₂	9	8	8	8	-	-	-	3	4	6	7
		SO ₂	2	-	1	1	1	1	1	1	1	1	1
75	Pomona-Walnut Valley	NO ₂	6	5	5	5	-	-	-	5	8	8	9
		SO ₂	1	-	2	1	1	2	2	-	2	2	2
76	South West Coastal	NO ₂	9	5	3	2	-	-	-	1	2	2	1
		SO ₂	4	-	2	2	2	1	2	2	2	2	1

* All times used are Pacific Standard Time.

APPENDIX D
WEATHER DATA FOR THE PERIOD MAY 21st TO MAY 24th, 1968

This appendix gives the following weather data for the Los Angeles Basin during the period of our investigation:

- (a) A brief summarizing statement of the daily weather.
- (b) The upper air temperature/pressure sounding at Los Angeles.
- (c) Surface weather for the following stations:-

LAX	Los Angeles Airport
BUR	Burbank
LGB	Long Beach
SAN	San Diego
SBA	Santa Barbara
SDB	Sandberg
SBD	San Bernardino
RIV	Riverside
BUO	Beaumont
PSP	Palm Springs
TRM	Thermal
IPL	Imperial
ONT	Ontario

These data were compiled by the Geography Department of the University of California, Riverside.

TUESDAY, May 21st, 1968

Stratus with a ceiling of about 1000 feet persisted over the coast all night and invaded the inland valleys at 4 a.m. At sunrise the stratus lifted (to a ceiling of about 3000 feet) and became quite unstable - a light shower fell at Riverside and others were observed in the Perris block. Broken cumulus to stratocumulus with a ceiling of about 3000 feet persisted the rest of the day. Visibilities were good at all areas all day and temperatures much cooler. The Coachella Valley again experienced dust in the evening hours. Soundings indicated an inversion at 4000-5000 feet and general cooler temperatures aloft as a result of an approaching trough.

WEDNESDAY, May 22nd, 1968

While scattered stratocumulus clouds occurred over coastal valleys in the early morning, surface heating produced a broken to complete stratocumulus cloud cover after sunrise, while the cloud cover persisted all day, some clearing took place along the immediate coast. Ceilings were about 3000 feet.

Soundings reported an inversion at 4000 feet which was largely destroyed by evening hours.

Visibilities were again good except in the San Bernardino -Riverside area in the early morning.

Deserts experienced severe dust storms most of the day and temperatures were again much cooler.

THURSDAY, May 23rd, 1968

A broken stratocumulus cover with a ceiling of 4000 feet resulting from morning surface heating, dominated skies over the coastal valleys, although more extensive clearing took place in the afternoon. Soundings indicated an ill defined inversion at about 5000 feet.

Persistent breezes maintained the good visibility pattern at all areas.

The air aloft remained relatively cool and slightly unstable, particularly over the interior (LAS VEGAS), as a trough passed into the Great Basin. As a result, flow aloft began to shift from westerly to northwesterly, a pattern which continued into Friday.

Dust storms were less severe in the Coachella Valley.

FRIDAY, May 24th, 1968

As the trough continued eastward from the Great Basin, the somewhat unsettled weather of previous days disappeared, temperatures warmed up, winds decreased, and visibilities also decreased. Cloudiness was restricted to a few cumulus in the morning hours and a band of cirris in the evening. There was considerable smoke and haze along coastal areas until noon and interior valleys until evening.

Sondings indicated redevelopment of an inversion at about 2000 feet.

Tuesday

May 24, 1968

HOUR Midnite 0000
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (in)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
WAX	Stratus Overcast 700 ft	8	1014.5	61	57	SW	8	ceiling rigid lower west
BUR	broken Stratus 2500 ft	15		63	58	SE	4	
LEG	Stratus Overcast ceiling 200 ft	3 fog	1019.2	60	58	S	7	
SAW	" 1000 ft	12	1015.2	65	57	NW	2	
BRA		10	1014.6	56	55	-	0	
SDS		-						
SDS		8		58	54	-	0	
BIV		7		56	50	-	0	
BUD								
POJ								
TRH		15	1007.5	80	52	N	15	
IRL		20	1008.5	80	50	NW	8	
ONT		10		60	56	SW	1	

Tuesday

DATE May 21, 1968

HOUR 10 am 1000
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (H3)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	broken stratus-cumulus 2500 ft	10	1014.0	65	55	SW	7	
BUR	"	10		64	55	SE	5	
LED	Stratus overcast 2500 ft 1500 ft	6 Haze Smoke	1014.6	65	57	N	4	
SAN	" 1900 ft	15	1014.9	64	53	SW	6	Stratocumulus NE-SE
SBA		15	1014.6	63	53	S	9	heavy cumulus topping mtns
SDB	broken sky 800 ft	20		45	36	N	17	
SSD	broken Stratus 3500 ft	15		64	50	-	0	
RIV	" 3000 ft.	30		64	48	S	6	
BUC								
PSP		30		82		NW	17	
TRM		10	1007.8	81	46	NW	12	Browning dust SE-SW
IDL		40	1007.8	89	26	W	15	
ONT	broken Stratus 2500 ft 1600 F	4 Haze		68	53	W	4	

Tuesday

DATE May 21, 1968

HOUR 11 am (100
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (in)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	broken Strato Cumulus 3000 ft	10	1014.5	66	55	SW	10	
BUR	"	10		67	53	SE	3	
LOS	" 2500 ft	7	1014.9	67	57	S	8	
SAH	Stratus Overcast 2700 ft 2000 ft	15	1014.9	65	52	SW	10	
SBA		20	1014.9	65	51	S	8	
SDB				47	36	N	16	
SBD	Few cumulus 5000 ft	25	1012.8	68	50	SE	5	
RIV	broken cumulus 3500 ft	30	1012.4	68	47	S	4	
BUO	Few Strato cumulus	25		65	48	SW	5	
PSP		30		83		NW	15	Gusts to 30
TRK		15	1007.5	85	39	N	5	
IDL		40	1007.8	92	30	W	17	Gusts to 24
ONT	broken Strato Cumulus	7		70	48	S	5	

DATE May 21, 1968 Tuesday

NOON
HOUR ~~11 AM~~ 1200
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (mb)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LOS	few cumulus 3700 ft	10	1014.4	68	55	SW	10	
MTJ	11 3500 ft 1500 ft	10		70	53	SE	10	
LEO	2500 ft 1500 ft	12	1014.6	69	56	S	11	
SAN	broken stratus 2200 ft 3000 ft	30	1014.9	66	51	SSW	7	
SEA		20	1014.9	67	51	SE	10	few cumulus along mtns
SDB	broken cumulus 1000 ft	50		51	36	NNW	10	strato cumulus tapping higher mtns
SJD	Few Cumulus 3000 ft			71	50	W	1	
RIV	11 3500 ft			70	47	SSE	2	
CLD								
PSP		30		85	55	NW	15	
TRN		15+	1007.1	88	39	N	7	
IDL			1007.1	95	31	W	15	6-8 sts to 22
JNT	broken strato Cumulus 2500 ft 1600 ft			72	52	SW	9	

DATE May 21, 1968 Tuesday

HOUR 1 pm 1300
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (in)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		10	1014.5	70	56	SW	13	Stratocumulus E-S
DBS	broken stratocumulus	10		73	54	SSE	6	
LGS	"	15	1014.6	69	56	SSW	8	
GAN	"	30	1014.9	70	52	SW	9	
SBA	Few cumulus 3500 ft	20	1014.9	67	51	SE	10	
SDB	broken sky ceiling 1500 ft	50		52	36	NNW	9	Stratocumulus hyalimms
SBD	Few cumulus 5000 ft	25		74	50	NW	6	
AIV		35		71	48	SSE	6	Few cumulus
SUG		25		67	46	W	3	
PSZ		30		88		NW	22	
TDM		20	1006.4	92	38	WNW	7	
EPL		40	1006.4	94	30	NW	15	Gusts to 21
ONT	Few cumulus	7	-	72	52	SW	10	

2175 May 21, 1968

Tuesday

HOUR 2 pm 1400
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (in)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		10	1014.0	71	56	SW	12	Status E-S
BUR	broken Strato Cumulus	15		73	54	SE	6	
LGB	"	20	1014.2	70	55	S	10	
SAW	Few cumulus 2700 Ft 3500 Ft	20	1014.6	61	50	SW	8	
SBA	"	20	1014.2	67	52	SE	10	
SCB		50		54	36	NNW	11	Strato Cumulus higher mts
SBD	Few cumulus 5000 Ft	15	1011.6	74	51	WNW	10	visibility lower w
RIV	" 3500 Ft	35	1011.4	72	48	SE	7	
BOB								
PSP		30		89		NW	30	
TRM		20	1005.8	92	37	NNE	16	
IPL		40	1006.4	96	37	WSW	20	Custs to 28
ONT	Few cumulus 3000 Ft	7		74	52	SW	17	

DATE May 21, 1968

T-214

HOUR

3pm 1500
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	Comments
LAX		15	1013.6	69	55	WSW	12	Stratocumulus ENE-S
BUR	Broken Strato-cumulus 4500 ft	15		72	53	S	5	
LGB	Few strato-cumulus 2500 ft	20	1013.9	72	55	SSE	11	
SAH	" 2700 ft	20	1014.2	68	50	SW	7	
SBA		25	1013.5	67	50	WSW	12	Gusts to 20 Cumulus N-NE
SDB				56	36	NNW	13	Stratocumulus along highway
SDD		8		73	51	NW	9	Cumulus N + S
SIV	Few cumulus 3500 ft	25		72	51	NW	9	
SUG		25		74	48	SW	6	
PSP		30		89		NW	30	
TRM		10	1005.1	93	35	NW	11	
IFL		40	1005.8	95	39	WSW	18	Gusts to 25
ONT	Few Cumulus 3000 ft	10		74	52	WSW	11	

DATE May 21, 1968HOUR 4 pm 1600
(Pacific Daylight Time)

STATION	Obs.	Visibility	Sea Level Pressure (10)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
222		15+	1013.1	70	53	WSW	15	
225	Broken Cumulus 4500 ft	15		71	53	SSE	4	
228		25	1013.2	70	55	SW	10	Few Stratocumulus, alto cum-lus
230	Few-Strato cumulus 3500 ft	20	1013.9	68	51	SW	10	
232		25	1012.5	67	49	WSW	18	cumulus dissipation
233				54	31	N	10	
235	Few cumulus 7000 ft	8		75	54	W	9	
237		20		73	51	NW	11	GUSTS 17 Stratocumulus N-E + S
240								
242								
244		15+	1004.4	95	39	N	8	
246		30	1004.7	94	32	W	19	GUSTS to 27
247	Few cumulus 3000 ft	10		72	52	SW	14	

DATE May 21, 1968HOUR 5pm 1700
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (in)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		20	1012.8	68	53	SW	12	
EUR	broken cumulus	20		69	52	S	4	
LGB		25	1013.2	70	53	W	12	
SAN	Stratus overcast 3000 ft	20	1013.9	65	53	SW	18	
SEA		25	1022.2	67	48	SW	15	Gusts 22. Few cumulus NW
SDB		8		55	37	N	11	
SBD	Few cumulus 7000 ft	15	1010.8	73	53	W	9	
RIV		25	1010.4	71	50	NW	11	Stratocumulus E-S
SND	Broken cumulus 6000 ft			72	49	SW	6	
PSP	-	15						
TRM		15 15	1004.1	93	45	N	15	Gusts 20. Dust S-W-N
IPL		30	1004.4	92	31	W	15	Gusts 25
ONT	Few cumulus 3000 ft	8		71	52	SW	14	

Wad
DATE May 22, 1968

HOUR Midnight
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		10	1012.5	58	52	SW	6	Few stratus, alto cumuli NE
BUR	Few cumulus 4000 ft	10		60	50	SE	3	
LGB		12	1013.2	59	56	S	7	Few stratus
SAW	Stratus Overcast 2300 ft	10	1013.5	63	53	W	8	
SBA		15	1012.9	61	42	SW	5	
SDB				44	32	NW	19	
SBD	Overcast 6000 ft	10		58	49	SW	3	
RIV	Few cumulus 6000 ft	15		57	48	SE	2	
SUD								
PSP								
TRM		7	1005.8	69	49	NW	20	Cusrs 25, dust clouds
IPL		20	1006.1	66	44	W	15	Cusrs 20
ONT	broken Cumulus 4000 ft	15+		59	49	SW		

DATE May 22, 1968HOUR 10 am
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (mb)	Temp (°C)	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	few stratus cumulus 1800 ft	10	1013.3	68	54	SW	5	
BUR	" 3000 ft	10		65	49	SE	6	
LOS	" 2000 ft	14	1013.9	65	54	S	6	
SAN	" 3500 ft	15	1014.2	66	50	S	8	
UOA		25	1013.9	61	44	S	6	
GGG		50 +		43	31	NNW	20	Rest of clouds N large cumulus NE
GGG	broken stratus cumulus 4500 ft	8		61	49	WNW	4	
RIV	" 2400 ft 5500 ft	20		62	45	W	3	
LOS	overcast 1500 ft	3 Haze		55	47	SW	5	
PSP	few high clouds	10		77		NW	6	clouds in Pass NW dust N, NE, E, SE
TRM	clear	7	1007.5	76	45	NW	16	dust, 22 DUST
IFL		30	1007.8	75	40	W	12	
ONT	broken stratus cumulus	9		61	49	WSW	7	

DATE May 22, 1968HOUR 11 am
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MD)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	few stratus cumulus 2000 ft	15	1013.5	69	54	WSW	10	
BUR	" 3000 ft	10		68	50	SE	5	
LGB	" 2000 ft	15	1014.2	66	54	SSE	8	
SAN	3500 ft 5000 ft	15	1014.2	67	49	SW	6	
SBA		25	1013.5	72	44	NW	7	
SDB		50+		46	32	NNW	16	large cumulus N
SSD	broken stratus cumulus 4500 ft	10	1012.7	63	47	W	6	
RIV	" 5000 ft	20	1012.0	63	44	W	2	
DUO	overcast 3000 ft	3 Haze		57	45	W	5	
PSP		10		79		NW	20	Dust E
TRK		7	1007.8	78	46	NW	20	Dust
IPL		30	1007.8	79	39	NW	12	Dust
JNT	broken stratus cumulus 1600 ft 3800 ft			63	49	ESE	7	

DATE May 22, 1968HOUR Noon
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	broken stratocumulus 2000 ft	15	1013.7	68	55	SW	13	
BUR	3000 ft	10		70	47	SSE	5	
LGB	2000 ft	15	1014.2	67	52	SW	8	
SAN	3500 ft	15	1014.6	67	50	SW	8	
SBA		25	1013.5	71	45	SW	15	Gusts to 20 few cumulus NE
SDB		50 +		49	34	N	14	large cumulus over Mtn tops
SBD	broken stratocumulus 5000 ft	15		65	47	NW	8	
RIV		20		66	44	WNW	7	
BUO	—							
PSP		10		79		NW	32	Dust NW-SE
TRM		6 Dn	1007.5	80	44	N	14	Gusts 22 Dust
IPL		30	1007.8	86	40	WNW	12	Gusts 19
ONT	broken stratocumulus 1500 ft 3800 ft			67	48	W	7	
AVNY	Broken cumulus 4500 ft	10				SSE	12	
FUL	broken cumulus 2000 ft	10				SSW	11	

DATE May 22, 1968HOUR 1 PM
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	few cumulus 2000 ft	15	1013.1	68	55	SW	10	
OUR								
LGB	2000 ft	15	1013.9	69	52	S	11	
SAN	3500 ft	15	1014.2	67	52	SW	7	
SBA		25	1012.5	69	46	SW	21	Gusts 28 Dist air few stratocumulus EN
SDB		50+		51	33	N	10	Mtn fogs observed by large cumulus
SBD	broken stratocumulus 5000 ft	15		69	48	WNW	7	-
RIV	"	12		68	44	NW	5	Visibility lower W-N
BUO	broken stratocumulus 5000 ft	5 Haze		64	44	SW	5	
PSP		10		81		NW	25	Gusts 35 Dust E
TRK		6 Dist	1006.4	83	41	NW	12	Gusts 18
EFL		30	1007.1	85	41	SW	15	Gusts 25
ONT	broken strato cumulus	9		68	55 48	SW	8	
WNY	Broken cumulus	10				SSE	20	
FUL	4500 ft							

DATE May 22, 1968HOUR 2 pm
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (HS)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	few Strato Cumulus 2000 ft	15	1012.8	68	55	SW	13	
DUR	" 3000 ft	10		72	50	S	7	
LSB	" 2000 ft	15	1013.5	69	52	SSW	12	
SAN	" 3500 ft	15	1013.9	68	51	SW	11	
SEA			1012.2	69	48	SW	15	Gusts 25
SDS		50 +		53	33	NW	13	Mtn tops obscured by large cumulus
SBO								
RIV								
BUC								
PSP								Dust E
TRH		6 Dust	1006.1	86	43	NW	14	Gusts 22 Dust
LPL		30	1006.5	87	41	SW	14	
DNT	broken cumulus 4000 ft	9		69	48	SW	8	
VNY	broken cumulus 4500 ft	10				SSW	15	
FUL	few cumulus 2000 ft	20				S	10	

DATE May 22, 1968HOUR 3 PM
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		15+	1012.6	69	51	SW	12	large cumulus NE-SE
BUR	few cumulus 3000 ft	10		72	48	S	6	
LGB	" 2000 ft	15	1013.2	70	52	SW	11	
SAN	" 2600 ft	15	1013.2	67	51	SW	9	
SBA		25	10115	72	46	W	20	Gusts 26 few stratocumulus tops N.
SDS		50+		56	34	N	7	Mtn tops obscured NW-NE
SBD	broken stratocumulus 5000 ft	15		68	47	W	7	
ONT	"	20		71	44	WNW	6	moderate cumulus E-S
BUG	broken strato cumulus	5 Haze		65	44	SW	4	large cumulus
PSP		10		83		NW	25	Gusts 35
TRM		7	1005.1	85	41	NNW	16	Dust
IPL		30	1006.1	86	41	WSW	14	Gusts 21
ONT	broken stratocumulus 4000 ft			69	49	SW	11	
VNY	broken cumulus 4500 ft	10				SSW	15	
FUL	few cumulus 2000 ft	10				SW	10	

DATE May 22, 1968

HOUR 4 PM
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		15+	1011.9	68	50	SW	20	Gr Cumulus E-SE
BUR	Few strato-cumulus 3000 ft	25		73	45	S	5	
LGB	"	14	1012.5	71	47	W	14	
SAN	2500 ft	15	1012.9	67	52	SW	12	
SEA		20	1011.2	72	46	W	20	Gusts 27 Few Strato cumulus NNE-EW
SDB				53	31	NNW	15	
SBD	broken Strato cumulus 5000 ft	12		67	49	W	10	
RIV	Few Strato cumulus 5000 ft	20		70	44	NW	12	Gusts 21 Moderate cumulus NE-SE
BUO	-							
PSP		10		82		NW	25	Gusts 35 Dust NW-SE
TRH		7	1004.4	85	42	NNW	16	Gusts 22
IPL		30	1005.8	85	42	W	13	Gusts 24
ONT	broken cumulus 4000 ft	7		67	50	WSW	17	

DATE May 22, 1968HOUR 5 pm
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		15+	1012.6	59	51	SW	12	large cumulus NE-SE
BUR	Few cumulus 3000 ft	10		72	48	S	6	
LEG		15	1013.2	70	52	SW	11	
SAN	11 2600 ft	15	1013.2	67	51	SW	9	-
SBA		25	1011.5	72	46	W	20	Cvsts 28 Stratocumulus mtns NE-NW
SDB								
SED	broken stratocumulus 5000 ft	15		68	47	W	7	
RIV	11	20		71	44	WNW	6	Moderate cumulus E-S
BVO	11	5 Haze						
PSP		10		83		NW	25	Cvsts 35 Dust E-NE
TRM		7	1005.1	85	41	MNW	16	Dust SW-MW
IPL		30	1006.1	86	41	WSW	14	Cvsts 21
DNT	broken stratocumulus 4000 ft			69	49	SW	11	
MVY	Few cumulus 4500 ft	15				SW	12	
FUL	Few cumulus 2000 ft	10				SSW	10	

Times

D 22

DATE

May 23

HOUR

~~Midnight~~ Midnight
(Pacific Daylight Time)

STATION	Clouds	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX		10	1011.5	58	52	S	5	
ORF		15+		59	50	ESE	4	
LGS		14	1012.5	58	51	S	5	
SAN	broken 5-10-5	15	1013.2	61	50	W	5	
SEA		15+	1012.5	59	45	NNW	15	
SEA		—						
ED		—						
ADV		15		51	44	SSE	2	
BLD								
REP								
TRM		8	1006.1	68	50	N	20	
PL		20	1007.1	63	41	W	13	6-23

JNT

15+

53

47

NW

8

thurs
DATE May 23, 1968

HOUR 10 am
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (HG)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	broken stratus 2000 ft	7	1013.9	66	48	ESE	10	
BUR	Few stratus 2500 ft	15		63	48	SE	4	
LGB	1800 ft	12	1014.6	64	51	S	7	
SAN		25	1014.2	65	49	SSW	12	few stratocumulus
SDA	cirrus RW	15	1014.6	62	50	SE	10	
SDS				49	31	NNW	7	Mtn tops obscured N
SBD	broken stratocumulus 5000 ft	10		59	46	-	0	
IV	2800 ft	20		58	44	-	0	
DUO	-							
FSP		20		74		NW	8	Dust E
TRM		6 Dust	1009.5	74	40	NNW	12	
EPL		40	1009.8	75	39	NW	10	
DNT	Broken stratus 2200 ft	7			48	SW	4	

DATE May 23, 1968HOUR 11 am
(Pacific Daylight Time)

STATION	Cloudiness	Visibility	Barometric Pressure (MB)	Temperature	Dew Point	Wind Direction	Wind Speed	COMMENTS
LAX	few stratus 2200 ft	8	1014.3	65	55	SW	9	
BUR	" 3000 ft	15	"	65	59	ESE	5	
LGB	" 2000 ft	15	1014.6	67	52	S	9	
SAH		20	1014.6	66	49	SSW	8	
SEA		15	1014.9	61	50	SSE	7	
SDS				52	31	N		
SBD								
RIV								
EVO	broken overcast 3500 ft	5 Haze		56	43	WSW	4	
PSP		30		77		NW	10	
TAM								
IPL		40	1009.5	79	39	NNW	6	
ONT	broken overcast 2200 ft	7		61	47	NW	7	
VNY	few	10				S	8	
FUL	few cumulus 2000 ft	7				S	10	

DATE May 23, 1968HOUR NOON
(Pacific Daylight Time)

STATION	Pressure	Visibility	Barometric Pressure (mb)	Temperature	Dew Point	Wind Direction	Wind Speed	Comments
LAX	few cumulus 2500 ft	10	1014.3	66	57	SW	9	
ONT	broken strato cumulus 2400 ft	10		69	48	S	5	
USE	" 2000 ft	15	1014.9	68	52	SSW	6	
SAN		20	1014.6	68	49	SW	8	
SEA		15	1014.9	62	50	SE	9	Few cumulus over mtns
SDB				52	31	N	11	
SRE	few cumulus 5000 ft	10		64	46	0	-	
RVV	" 4000 ft	20		60	43	NW	9	
EUO								
PCS		30		81		NW	15	
ELI								
DEL		30	1009.5	81	38	W	8	
ONT	broken strato cumulus 2000 ft	6 Haze		64	48	N	7	
VNY		10				S	8	
FUL	few cumulus 2500 ft	7				SW	10	

Thurs
May 23, 1968

1 PM

	few cumulus 2500 ft	12	1014.0	67	57	SW	12	
	2000 ft	15	1014.6	66	51	SSE	7	
		20	1014.6	68	49	WSW	9	
		15	1014.9	65	47	E	9	
		50		52	31	N	14	
	few cumulus 5000 ft	10		64	45	W	4	
	4000 ft	15		67	43	NW	10	
		25		65	43	SW	3	cumulus clouds
		30		82		NW	22	
		15	1008.8	83	35	E	4	
		50	1009.5	84	30	WNW	7	
DNT	broken cumulus 2500 ft	6 Haze		69	47	NW	1	
VNY		10				SW	8	
FVL	few cumulus 2000 ft	7				SW	10	

May 23, 1968

2 PM

	Few cumulus 2500ft	13	1014.0	67	55	SW	15
	broken cumulus 2500ft	8		70	50	S	7
	"	15	1014.2	69	51	S	12
200		20	1014.2	68	51	SW	10
200		20	1014.2	65	51	SE	12
200		50		55	32	NNW	17
200		10	1012.9	67	45	NNW	6
200	Few cumulus 4000ft	10	1012.1	70	43	NNW	5
200	—						
200		25		83		NW	23
200		15+	1007.8	84	36	NNW	8
200		50	1008.1	88	37	NNW	8
200	Few cumulus 2500ft	7		72	46	NNW	1

Few cumulus
MINS

Stratocumulus
S

NOT REPRODUCIBLE

May 23, 1968

3 PM

	Direction	Temp	Hum	Wind Dir	Wind Spd	Remarks
	12	1013.8	66	57	SW	16
						Few cumulus Stratocumulus
few strato-cumulus 2500 ft	8		72	49	ESE	8
few cumulus 2500 ft	15	1013.9	70	51	S	11
	20	1014.2	68	52	SW	8
	20	1014.2	65	52	SE	10
	50		55	32	NNW	12
	10	6	69	44	N	7
						Stratocumulus S-SW
few cumulus	15		72	43	NW	8
	25		83	50	NW	23
						Est 33
	15+	1007.5	86	35	SW	6
	40	1007.5	90	40	N	3
ONT	cumulus 2500 ft	7	72	47	N	1
VNY						
FUL	few cumulus 2500 ft	7				8
VNY	broken cumulus 3000 ft	15				10
FUL	2500 cumulus	15+			SE	10
					S	10

NOT REPRODUCIBLE

May 23, 1968

4 PM

		Pressure	Temp	Humidity	Wind Dir	Wind Spd
few cumulus 3000 ft ↘ "	10		72	49	SE	8
	20	1013.5	71	48	S	11
	20	1013.9	67	52	SW	7
	20	1013.2	71	48	SW	11
few cumulus 5000 ft "	10		72	46	W	10
	15		72	44	NW	8
	20		85		NW	22
	15+	1007.1	87	35	N	10
	40	1006.8	91	36	SW	10
JNT	few cumulus 2500 ft	7	72	48	NW	
VNY	Broken cumulus 3000 ft	15			SE	10
FUZ	2500 few cumulus	15+			S	10

NOT REPRODUCIBLE

May 23, 1968

1300 5 pm

		14	1013.3	66	51	WSW	14 Few cumulus S
	Few cumulus 3000 ft	15		72	48	SW	6
		20	1013.5	69	48	WNW	9
	Few stratus cumulus 1800 ft	20	1013.9	66	52	WSW	6
		20	1012.5	74	46	MNW	8 smoke layer off shore few cumulus mths ↑
				56	30	N	16
	Few cumulus 5000 ft	10	1012.0	71	48	WSW	10
	" 4000 ft	20	1011.3	70	46	NW	12
	3 Haze			70	41	W	5 Cumulus SE-S
	20			83		NW	18 Gusts 30
	15+	1006.1	57	37	NW	9	
	30	1006.4	90	36	W	15	Gusts 23
ONT		7		70	49		
RUZ	Few cumulus 2500 ft	15+				S	10
VNY	" 3000 ft	15				S	15

May 23 Thurs

11 PM

	12	1013.6	57	50	WSW	8
	10		60	49	SE	3
	15+	1014.2	58	50	N	4
broken stratus 1900 ft	15	1014.6	61	51	SW	5
	15+	1014.2	59	45	NW	12
			45	28	NNW	23
broken stratocumulus 5500 ft	8	1013.6	57	47	—	0
Few cumulus 4500 ft	10	1013.3	55	43	—	0
—						
—						
	10	1007.8	70	47	NNW	18
	20	1008.5	68	35	W	13
ONT	15		57	46	21	

NOT REPRODUCIBLE

May 25

9 am

Few cumulus 2000 ft	C Haze Smoke	1016.0	64	52	SE	6
	12	1016.0	62	50	SE	3
broken stratus- cumulus 2000 ft	8	1016.6	65	53	SE	7
	20	1016.3	64	48	SSW	5
	15+	1016.6	62	51	SE	9 Hazy east
	50+		49	28	NNW	11
Few cumulus 4000 ft broken cirrus	C Haze Smoke		57	48	—	0
broken stratus 1000 ft	7		54	45	S	4
Few cumulus 1500 ft	2 Haze		53	46	NW	4
	15		72		NW	16
	7	1012.2	72	45	NNW	9
	30	1012.2	72	42	NW	7
ONT Few Cirrus	5 Haze		59	49	—	—

F 81
May 24, 1968

NOT REPRODUCIBLE

10 am

Time	Wind	Temp	Humidity	Pressure	Clouds	Remarks
0900	S	67	52	1016.4	9	few cumulus 2000 ft
1000	ENE	65	47		10	
1100	SSW	67	53	1016.9	11	
1200	SSW	65	49	1016.3	20	
1300	SE	64	52	1016.9	15	
1400	NNW	51	29		50+	
1500	-	58	48		6	broken stratocumulus 4000 ft Haze Smoke
1600	S	58	45		7	few cumulus 1500 ft
1700	W	56	46		3	Haze
1800	NW	70			15	
1900	NW	75 77	42 41	1012.2	27	High clouds
2000	NW	77	41	1012.5	27 30	
2100		63	49		4	Few cirrus Haze

Fri May 24, 1968

11 am

		10	1016.6	68	53	SW	10	Cumulus N-E-
		10		70	50	S	5	
	Few cumulus 4000 ft	10	1017.3	68	54	S	9	
		20	1016.3	68	49	SW	8	
		20	1017.3	65	51	E	10	
		50+		54	22	NNW	12	
	Few cumulus 4000 ft	6 Haze Smoke	1016.3	61	49	-	0	
		7	1015.9	65	45	ENE	2	
		5 Haze		61	43	W	4	
		15		83		NW	4	
		10	1012.2	80	40	-	0	
		20	1012.5	80	42	NW	8	
JNT	Few cirrus	4 Haze		67	48	SW		

NOT REPRODUCIBLE

Fri May 24, 1968

Noon

	Pressure	Temp	Humidity	Wind Dir	Wind Speed	Remarks
AS	12	1016.9	68	53	SW	12 Cumulus N-E-S
BS	7		71	50	SE	5
CS	10	1017.3	62	53	S	8 few cumulus 2500 ft
DS	20	1016.9	68	50	SW	10
ES	20	1017.3	63	50	SE	10
FS	50 +		57	25	N	- Cirrus N
GS	6 Haz Smoke		66	50	-	6
HS	10		67	44	WSW	2
IS	5					
JS	15		83		E	9
KS	15	1012.2	85	39	NW	5
LS	20	1012.5	82	39	N	6
ONT	4 Haz		71	49	SW	

NOT REPRODUCIBLE

Fri
May 24, 1968

1 PM

12	1016.9	69	52	SW	12	- few cumulus	
7		73	51	SE	5		
14	1017.3	70	52	S	10		
20	1016.9	69	50	W	9		
20	few cumulus	1017.6	62	51	SSE	8	
—	50+	61	30	N	10		
7		70	48	W	6		
10		70	43	WSW	5	Gusts 11	
5 Haze		68	43	SW	4		
20		86		E	3		
15	1011.5	88	34	SE	6		
20	1012.2	85	36	NW	8		

ONT

VNY

FVZ

few cumulus

few cumulus

2500 ft

1.0

7

S

SSW

14

4

NOT REPRODUCIBLE

May 24, 1968

2 pm

RECEIVED MAY 27 1968

		Pressure	Temp	Wind	Wind	Wind	Wind
				Point	Speed	Dir	Speed
103	Few cirrus	12	1016.6	69	52	SW	12
104		7		74	52	SSE	7
105	Few cirrus	15	1016.9	72	52	SW	12
106		20	1016.9	68	50	NW	5
107	Few cirrus	20	1017.3	65	49	SE	10
108	"	50 +		59	30	N	11
109	6 Haze Smoke	1015.3	72	46	S	G	
110		7	1015.2	71	43	WNW	10 Gusts to 20
111							
112		20		87		NW	10
113		15	1011.2	89	31	E	8
114		20	1011.9	88	42	NNE	7
ONT	Few cirrus	5 Haze		75	47	W	10
VNY	"	10				S	15
FUL		5 Haze				SSW	8

Fri May 24, 1968

3 pm

		Temp	Bar	Wind	Wind Dir	Wind Spd	Clouds
	Few cirris	13	1016.6	69	53	SW	13
	"	7		75	52	S	5
	"	15	1016.9	73	50	SSW	13
		20	1016.6	67	51	W	9
		20	1017.3	63	52	SE	12
		50+		59	31	N	23
	Few cirris	7		76	47	W	5
		3 Haze		75	44	W	6
		20		90		NW	11
		15+	1010.8	90	29	N	6
		25	1010.5	88	36	NW	9
ONT	Few cirris	4 Haze		77	48	SW	11
VMY	"	10				SE	14
FVL	"	5 Haze				SW	9

May 24, 1968. Fri

4 PM

	Clouds	Barometric Pressure	Temperature	Dew Point	Wind Direction	Wind Speed
SEA	few cirrus	10	1016.4	68	54	SW 11
SEA	↖	7		75	51	S 4
SEA	↖ " "	15	1016.3	75	49	NW 10
SEA	" "	20	1016.3	70	50	NW 9
SEA		20	1016.9	63	51	SE 8
SEA				60	31	NNW 21
SEA	broken cirrus	7		76	47	WSW 6
SEA	" "	8		75	45	NW 10
SEA	↖ -					
SEA	↖ " "	20		91	32	NW 19
SEA	few cirrus	15+	1011.5	93	30	E 6
SEA	↖	30	1010.8	88	36	N 8
ONT	↖ " "	4 Haze		76	49	SW 11
VNY	↖	8				SE 12
FUL	broken cirrus	10				S 10

Fr, May 24, 1968

5 PM

NOT REPRODUCIBLE

		Visibility	Barometric Pressure	Temperature	Dew Point	Wind Dir	Wind Speed	Comments
		8	1016.0	68	55	SW	11	
		20		73	49	S	9	
		15	1015.9	75	48	WNW	9	
		20	1016.3	69	52	NW	13	
		20	1016.3	65	49	SE	8	
				58	28	N	19	
ONT	overcast cirrus	6 Haze smoke	1014.4	75	49	W	13	increasing Haze smoke
UNY	broken cirrus	10	1014.5	74	44	NW	10	visibility higher S
FUL		3 Haze		74	43	NW	7	
	broken cirrus	20		83		NW	22	
	Few cirrus	15+	1009.8	91	32	-	0	
		30	1010.2	93	34	NW	7	
ONT		5 Haze		74	50	SW	11	
UNY		10				SE	12	
FUL		10				S	7	

Friday May 24, 1968

HOOR


6 PM

Wacific Day 30 1 1968

NOT REPRODUCIBLE

Time	Cloud	Visibility	Barometric Pressure (mb)	Temp (°C)	Dew Point	Wind Dir or Line	Wind Speed	Comments
1800		8	1015.8	67	54	SW	10	Few cirrus N
1815		20		72	49	S	5	
1830		15	1015.9	72	48	W	12	
1845		20	1015.9	67	53	NW	11	
1900	Few cirrus	20	1016.3	65	49	SE	7	
1915				56	27	NW	20	
1930	broken cirrus	6 Haze smoke		73	50	WSW	8	
1945	Few cirrus	10		72	44	NW	8	Gusts 15
2000	—							
2015	↙ "	15		81		NW	21	
2030	"	15+	1009.8	93	33	W	5	
2045		30	1009.8	90	35	N	7	


ONT



LAX


May 21, 1951


21835Z 0600 PDT Tuesday



LAX May 22 0600 PDT MORNING Wednesday

D 44





LA INVERSION
 DATA FOR MAY 24, 1968
 0600 P.
 BASE 2300 FT Temp 11.6 RH 54
 TOP 2720 Temp 13.5 RH 40

D 45

LAX May 24 24234 (m 50) 0600 PDT FRIDAY